

## ORIGINAL ARTICLE

WILEY

# Measured concentrations of consumer product chemicals in California house dust: Implications for sources, exposure, and toxicity potential

Hyeong-Moo Shin<sup>1</sup> | Christoph Moschet<sup>2</sup> | Thomas M. Young<sup>2</sup> | Deborah H. Bennett<sup>3</sup>

<sup>1</sup>Department of Earth and Environmental Sciences, University of Texas at Arlington, Arlington, Texas

<sup>2</sup>Department of Civil and Environmental Engineering, University of California, Davis, Davis, California

<sup>3</sup>Department of Public Health Sciences, University of California, Davis, Davis, California

**Correspondence**

Hyeong-Moo Shin, Department of Earth and Environmental Sciences, University of Texas at Arlington, 500 Yates Street, Box 19049, Arlington, TX 76019.  
Email: hyeongmoo.shin@uta.edu

**Funding information**

National Institute of Environmental Health Sciences, Grant/Award Number: P42-ES004699; U.S. Environmental Protection Agency, Grant/Award Number: EPA-G2013-STAR-K1

**Abstract**

Household dust is a reservoir of various consumer product chemicals. Thus, characterizing comprehensive chemical profiles of house dust may help improve our understanding of residential chemical exposure. We have previously developed a method for detecting a broad spectrum of chemicals in dust by applying a combination of target, suspect screening, and non-target methods with mass spectrometry preceded by liquid chromatography and gas chromatography. Building upon a previous study that detected 271 compounds in 38 dust samples, we presented concentrations of 144 compounds that were confirmed and quantified by standards in the same set of samples. Ten compounds were measured with median concentrations greater than 10 000 ng/g of dust: cis-hexadec-6-enoic acid, squalene, cholesterol, vitamin E, bis(2-ethylhexyl) phthalate, dioctyl terephthalate, linoleic acid, tricaprylin, tris(1-chloroisopropyl) phosphate, and oxybenzone. We also reviewed in vitro toxicity screening data to identify compounds that were not previously detected in indoor dust but have potential for adverse health effects. Among 119 newly detected compounds, 13 had endocrine-disrupting potential and 7 had neurotoxic potential. Toxicity screening data were not available for eight biocides, which may adversely affect health. Our results strive to provide more comprehensive chemical profiles of house dust and identified information gaps for future health studies.

**KEYWORDS**

concentration, dust, in vitro bioactivity assays, non-target, suspect screening, target

## 1 | INTRODUCTION

Thousands of chemicals are currently used in consumer products. Many of the consumer product chemicals have not been studied for exposure potential in the indoor environment where people in developed countries spend most of their time.<sup>1</sup> The most studied chemical classes in the indoor environment include pesticides, flame retardants, plasticizers, polycyclic aromatic hydrocarbons (PAHs), and per- and polyfluoroalkyl substances (PFAS).<sup>2,3</sup> Because of potential health concerns over exposures to some of these chemical classes, a large number of alternative chemicals are being introduced

into consumer products every year, following legislative activities or advocacy campaigns.<sup>4-6</sup> However, exposure and toxicity information needed to evaluate potential human health effects are limited for the alternative chemicals and other chemicals that were not previously measured in indoor environmental media.<sup>7</sup>

Chemical concentrations in indoor environmental media including air, airborne particles, and settled floor dust have been used to characterize residential exposure to indoor contaminants.<sup>2,8-10</sup> Many consumer product chemicals of current and emerging health concerns are semivolatile organic compounds (SVOCs).<sup>2</sup> When released from their original sources, SVOCs are

redistributed over time and primarily partitioned to dust and other indoor surfaces.<sup>3,11,12</sup> When dust concentrations are known but other media concentrations are not measured, partitioning models among dust, gas-phase, and airborne particles can be used to characterize residential chemical exposure.<sup>2,10</sup> Thus, there have been growing efforts in detecting and quantifying SVOCs in house dust.<sup>3</sup> However, a complete picture of the chemical fingerprint of dust (ie, identity and quantity of all chemicals present) is missing, because most previous studies analyzed *known* chemical classes via a targeted analytical method.<sup>2,13–17</sup> Therefore, development of advanced environmental monitoring methods has emerged as a prominent topic in indoor environmental research in order to detect both *known* chemical classes and those that were previously not targeted for detection in dust.

Advances in high-resolution mass spectrometry make it possible not only to detect *known* compounds for which reference standards are available (targets), but also to detect *expected* compounds using existing databases, libraries, or software matching algorithms (suspects) and even to identify previously *unknown* compounds (non-targets) through careful examination of high-resolution mass spectra.<sup>18,19</sup> To date, four studies have applied suspect screening and non-target methods to dust samples. Hilton et al<sup>20</sup> first applied a non-target method to one household dust sample obtained from the National Institute of Standards and Technology (NIST) using two-dimensional gas chromatography (GC × GC)-time-of-flight mass spectrometry (TOF/MS). Ouyang et al<sup>21</sup> carried out a non-target analysis for one household dust sample collected in Sweden using two-dimensional liquid chromatography (LC × LC)-TOF/MS. Rager et al<sup>22</sup> applied suspect screening and non-target methods to 50 household dust samples collected in the US from 2005 to 2006 using LC-TOF/MS. A comparative study of a non-target analysis was conducted in a composite house dust sample as part of a collaborative effort using LC-MS and GC-MS.<sup>23</sup> The methods used in the four studies are useful for identifying previously unknown and even unexpected chemicals in dust, but none of them presented concentrations that were confirmed and quantified by standards.

As part of an effort to evaluate a large number of environmental chemicals for potential health effects efficiently, the US Environmental Protection Agency (EPA)'s Toxicity Forecaster (ToxCast) program utilizes hundreds of *in vitro* high-throughput screening (HTS) assays to support prediction of *in vivo* toxicities.<sup>24</sup> In a parallel effort to screen a larger number of chemicals based on exposure, several high-throughput (HT) methods including exposure models have been developed to characterize and estimate exposures.<sup>10,25–28</sup> As indoor dust is a reservoir for SVOCs released indoors and can provide reasonable surrogates for characterizing exposures, we have previously developed a method for detecting a broad spectrum of chemicals in dust by applying a combination of target, suspect screening, and non-target methods using both LC-quadrupole time-of-flight (QTOF)/MS and GC-QTOF/MS.<sup>18</sup> Building upon this previous publication,<sup>18</sup> this current study presents chemical concentrations that were quantified in the same set of dust samples for target, suspect, and selected non-target compounds. In addition, we

### Practical Implications

- Our dust samples contain chemicals from various consumer product uses, including cleaning and personal care products, furniture, plastics, and pesticides.
- This supports the idea that dust can serve as a marker of use.
- We expect that this comprehensive investigation of chemicals present in dust will form the basis for future work to develop new hypotheses of adverse health effects due to exposures to previously overlooked compounds.

investigated whether the compounds detected in our samples had either endocrine-disrupting or neurotoxic potential and discussed possible applications of our findings to future health studies.

## 2 | MATERIALS AND METHODS

### 2.1 | Overview and scope of this study

The aim of this study was to inform key data gaps for assessing potential health effects for consumer product chemicals by integrating our measured dust concentrations with existing exposure and toxicity potential data. Our previously published study comprehensively characterized compounds found in house dust samples, detecting diverse and numerous consumer product chemicals. The present study extends those findings by quantifying concentrations, assessing household level variability, and considering potential exposure and toxicity of the compounds detected. Five steps were taken toward achieving the overall aim. First, we classified detected chemicals by their chemical class (eg, phthalate) and their common use category (eg, plasticizer). Second, we compiled information on chemical analysis techniques used to detect the compounds, including analytical instrument (LC or GC) and method (target, suspect screening, or non-target), the limit of detection (LOD), and information on whether identities were confirmed and concentrations were quantified by standards. Third, we summarized results from the chemical analysis, including the number of samples in which a compound was detected, information on whether a compound was newly detected in our house dust, and summary statistics of measured concentrations. Fourth, we added information on whether each detected compound has endocrine-disrupting or neurotoxic potential based on *in vitro* HTS assays. Fifth, we indicated whether the compounds detected in our dust have been biomonitoring in the US National Health and Nutrition Examination Survey (NHANES).<sup>29</sup>

Note that we did not compare our measured concentrations to those reported in other peer-reviewed studies but focused on summarizing our concentrations by chemical class or use category. Other studies have already ascertained that differences in concentrations among studies may result from different sampling methods as well as geographic and temporal variation in chemical use.<sup>2,30</sup> In

addition, we did not describe sample collection, dust extraction, and analytical methods in detail in the current study because the details are available in the previous publication.<sup>18</sup> The whole analytical method and workflow were completely validated, results of quality assurance (QA) and quality control (QC) for each of the analytical approaches were provided, and strengths and weaknesses of the various approaches and analytical instruments were discussed previously.<sup>18</sup> Thus, we briefly described sampling and analytical methods to the extent necessary for others to quickly extract key information regarding environmental monitoring, such as sample size, sampling method, period and location, and type of analytical instruments. Results of *in vitro* HTS assays presented in this study do not necessarily represent *in vivo* toxicities. Factors influencing toxicity such as pharmacokinetics and metabolism, early-life susceptibility, and genetic variability are not addressed by ToxCast.<sup>24,31</sup> Thus, toxicity potentials presented in this study need to be interpreted with caution. Other limitations of using *in vitro* HTS assays for predicting *in vivo* response are discussed elsewhere.<sup>31</sup>

## 2.2 | Sampling and analytical methods

We recruited 38 families in Northern California from May 2015 to August 2016. From each household, we collected one dust sample from an approximate 2 m<sup>2</sup> area in the main living room using a high-volume small surface sampler (HVS3), following a standard protocol.<sup>32</sup> Dust samples were sieved, and 100 mg aliquots were sonication-extracted with hexane/acetone (3:1 v/v) and acetone (100%). The extracts were then analyzed by both LC-QTOF/MS and GC-QTOF/MS with methods that were able to analyze compounds from various compound classes with widely differing chemical properties (eg, molecular size, logP). In addition to the classical target analysis using reference standards and isotope-labeled internal standards, additional suspect screening and non-target analysis were performed. In order to unambiguously confirm the identity of suspected and non-targeted compounds, additional reference standards were purchased if they were available. Details of quantification methods are available elsewhere.<sup>18</sup>

## 2.3 | Selection of target compounds

For the targeted method, we selected 76 chemicals for GC analysis and 56 chemicals for LC analysis (see Supporting Information [SI], Table S1). The targeted compounds included personal care products (PCPs; antimicrobial compounds, fragrance ingredients, parabens, and ultraviolet [UV] filters), markers of human inputs (skin oils and metabolites), flame retardants (polybrominated diphenyl ethers [PBDEs], organophosphate flame retardants [OP-FRs], and other FRs), pesticides (insecticides, fungicides, and herbicides), and a variety of other compounds widely detected in homes (phenols, phthalates, other plasticizers, PAHs, PFAS, and surfactants). The selection criteria included one (or multiple) indicator compounds from substance classes identified in our previous study<sup>7</sup> or compounds present in products listed in the US EPA's Consumer Product Chemical Profiles database (CPCPdb).<sup>33</sup>


## 2.4 | Chemical use categorization

Chemicals identified in the current study were classified into the most common and primary use category to understand the distribution of measured compounds by use category. For most compounds, we relied on "product" or "use" categorization available in databases such as CPCPdb and the US National Library of Medicine's Household Product Database (<https://householdproducts.nlm.nih.gov/index.htm>) to find the most common and primary use category of the compounds associated with at least one consumer product. For compounds with multiple uses, we also relied on web searches to find common uses. Multiple-use compounds were assigned to a primary use category and their secondary or tertiary use categories were further discussed in Results and Discussion. Thus, use categorization may be imprecise. For compounds that are consumed via dietary sources and also formulated in cosmetic products (eg, linoleic acid, palmitic acid, cholesterol, and fatty acids),<sup>34</sup> it is likely that emissions from cooking are the dominant source to residential floor dust. Therefore, we preferentially assigned their primary use category to food sources. We further discuss this in Results and Discussion.

## 2.5 | Data sources of endocrine-disrupting potential or neurotoxic potential

Many chemicals present in consumer products exhibit endocrine-disrupting potential<sup>35</sup> or neurotoxic potential.<sup>36</sup> To determine whether the compounds detected in our samples have either endocrine-disrupting or neurotoxic potential, we used *in vitro* HTS assays, most of which are included in the US EPA's ToxCast program. For endocrine-disrupting potential, we evaluated four main processes, including androgen, estrogen, thyroid, and steroidogenic. For androgen, we utilized androgen receptor (AR) pathway activity integrated from 11 AR-related *in vitro* HTS ToxCast assays and considered compounds with area under the curve (AUC) of  $\geq 0.1$  to be active in at least one AR pathway assay (active, inactive).<sup>37</sup> For estrogen, we utilized estrogen receptor (ER) interaction scores integrated from 13 ER-related *in vitro* ToxCast assays and considered compounds with an AUC score of  $\geq 0.1$  to be active in at least one ER pathway assay (active, inactive).<sup>38</sup> For thyroid, we utilized the results from the *in vitro* Amplex UltraRed-thyroperoxidase or thyroid peroxidase (AUR-TPO) assay<sup>39</sup> and a thyroid-specific *in vitro* HTS ToxCast assay.<sup>40</sup> Because decreased TPO activity reduces thyroid hormone synthesis, compounds that elicited a  $\geq 20\%$  reduction in maximal TPO activity were considered to inhibit TPO (active, inactive).<sup>39</sup> We also identified compounds that exhibited thyroid receptor activity measured by the *in vitro* ToxCast assay (active and inactive).<sup>40</sup> For steroidogenesis, we utilized results from a method that considered 10 steroid hormones, including progestogens, glucocorticoids, androgens, and estrogens using an *in vitro* HTS assay with H295R human adrenocortical carcinoma cells.<sup>41</sup> Among 2060 evaluated compounds, we considered compounds that altered at least 4 steroid hormones at the maximum tolerated concentration to be active (or inactive), the same criteria used in Karmaus et al.<sup>41</sup> For a neurotoxic indicator, we utilized microelectrode array hits as a measure of neural network activity in

**FIGURE 1** Summary of median concentrations (ng/g of dust) for 87 compounds (target + suspect + non-target) detected in more than 50% of the samples



vitro (yes, no).<sup>42</sup> A summary of toxicological endpoint data is provided in Table S2.

## 2.6 | Statistical analysis


Statistical analyses were performed using Microsoft Excel 2018. For concentrations between the limit of quantification (LOQ) and the LOD, we assigned a value of the LOQ divided by 2. For concentrations below the LOD, we assigned a value of the LOD divided by the square root of 2.<sup>43</sup> For compounds detected in more than 50% of the samples, we summarized measured median concentrations by five levels (<500, 500-1000, 1000-5000, 5000-10 000, and >10 000 ng/g of dust) to investigate which compound classes were measured and present at high concentrations. We also computed coefficients of variation (CV) to examine the variability of concentrations in dust across homes.

## 3 | RESULTS

### 3.1 | Measured dust concentrations

A total of 276 compounds were detected in our dust samples in which 5 additional compounds were detected after our previous study was published.<sup>18</sup> For 14 compounds, identification was not possible and only molecular formula (eg, C<sub>4</sub>H<sub>7</sub>FO) could be assigned. Table S2 summarizes information of analytical methods, results from the analysis, exposure, and toxicity potential for 262 detected compounds that could be identified with structure and

formula. Additional summary for all 262 compounds detected in our dust is provided in the Supporting Information (see Data S1 for overall description and Table S3 for summary by chemical class and analytical instruments/methods). Overall, a large number of UV filters, phthalates, and OP-FRs were detected in our dust samples and median concentrations for some of them were above 10 000 ng/g of dust (Figure 1). Ten compounds were measured with median concentrations greater than 10 000 ng/g of dust: cis-hexadec-6-enoic acid, squalene, cholesterol, vitamin E, linoleic acid, tricaprylin, bis(2-ethylhexyl) phthalate [DEHP], dioctyl terephthalate [DOTP], tris(1-chloroisopropyl) phosphate [TCIPP], and one UV filter (oxybenzone). Cis-hexadec-6-enoic acid, squalene, cholesterol, and vitamin E are found in skin surface lipids.<sup>44</sup> Linoleic acid, cholesterol, and tricaprylin are widely used in cosmetics and PCPs. However, it is likely that emissions from cooking may significantly contribute to the measured dust levels of linoleic acid, cholesterol, and vitamin E.<sup>34</sup> Consumer products (eg, electronics, plastic products, and shower curtains), building materials (eg, vinyl flooring), and furniture (eg, couches) are well-known emission sources of DEHP, DOTP, or TCIPP in the indoor environment. High concentrations of other chemical classes (eg, skin oils, cosmetic ingredients, and UV filters) detected in the current study highlight that humans and their activities, and possibly pets, play a role as sources of SVOCs in the indoor environment. Fungicides, PBDEs, PFAS, and pharmaceuticals were also abundant in our samples, but most were measured at concentrations below 500 ng/g of dust.



**FIGURE 2** Distributions of dust concentrations (ng/g of dust) for 56 target compounds detected in more than 50% of the samples. Notes: Two skin oils (sapienic acid and squalene) and linoleic acid (used in cosmetics and emitted during cooking) were grouped into “cosmetics” in this figure. Compounds with asterisk (\*) indicate the first measurement in household dust. This figure excludes two phenols (tetrachlorophenols and cresol) that were detected in most samples (above LOD) but that were below LOQ. See Table 1 for identification of abbreviations

In the present study, 119 compounds were identified and/or quantified for the first time in household dust (Figure S1). Some of these compounds were previously measured in US wastewater samples via target analysis but had not been measured in indoor dust. The majority of these compounds was detected via LC non-target (45%) and LC suspect (31%) approaches (see inset of Figure S1). These newly measured compounds mainly comprised surfactants ( $n = 25$ ), pharmaceuticals ( $n = 19$ ), compounds with unknown use information ( $n = 17$ ), and human metabolites ( $n = 12$ ), because of the polarity of these compounds. We also identified 6 phenols (some are also used as biocides) and 11 biocides (4 insecticides and 7 fungicides) in dust for the first time mostly via GC target and/or LC target analyses.

Overall, dust concentrations varied by almost three orders of magnitude across household samples and by almost four orders of magnitude across compounds (Figure 2). PFAS were measured at the lowest concentrations and had relatively large variability in concentrations. DEHP was shown to have the smallest variability (coefficient of variation,  $CV = 0.35$ ) across the samples. Except for tri-*n*-butyl phosphate (TNBP), OP-FRs were measured at higher concentrations than PBDEs, and bisphenol S (BPS) was measured at higher concentrations than bisphenol A (BPA). This is consistent with

recent changes in consumer use due to changes in product formulation and regulations affecting PBDEs and BPA. Variability metrics for all compounds including CVs are available in Table S2.

Below, we summarized our measured dust concentrations along with other exposure and toxicity potential information by four categories: (a) chemical classes other than biocides that have received considerable public attention in indoor dust (eg, phthalates, PBDEs, OP-FRs, and PFAS); (b) biocides (eg, insecticides and fungicides); (c) compounds in PCPs (eg, fragrance ingredients and UV filters); and (d) chemical classes whose dust concentrations are of less concern for environmental exposure calculations (eg, food additives and skin oils).

## 3.2 | Chemical classes of current and emerging concerns in indoor dust

### 3.2.1 | Phthalates and other plasticizers

Among 7 target phthalates, benzyl butyl phthalate (BBP), DEHP, di-isobutyl phthalate (DiBP), and di-*n*-butyl phthalate (DnBP) were detected in all of our samples. Median concentrations of these four phthalates were above 3000 ng/g of dust (Table 1). Diethyl phthalate

**TABLE 1** Summary of chemical classes that were detected in our dust samples (n = 38) and have been receiving considerable public attention

Chemical class	Compound name	Abbreviation	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
Phthalates	Benzyl butyl phthalate	BBP	G	T	50	1	38		9181	134 764	1	1	1
	Bis(2-ethylhexyl) phthalate	DEHP	G	T	50	1	38		39 124	77 532	0	0	1
	Di-isobutyl phthalate	DiBP	G	T	100	1	38		3465	25 917	0	1	1
	Di-n-butyl phthalate	DnBP	G	T	100	1	38		4974	15 983	1	1	1
	Diethyl phthalate	DEP	G	T	500	1	30		966	4184	0	0	1
	Dimethyl phthalate	DMP	G	T	50	1	27		103	388	0	0	1
Other plasticizers	Bis(2-ethylhexyl) adipate	DEHA	G	T	5000	1	15			17 522	0	0	
	Acetyl tributyl citrate	ATBC	G	T	50	1	38		7969	27 543	0	0	
	Diocetyl terephthalate	DOTP	G	T	50	1	38	1	35 678	115 606	0	0	
	1,3-Diphenylguanidine		L	T	30	1	38	1	3218	9659	1	1	
	Toluene-2-sulfonamide		L	S <sup>h</sup>	<100	1	38	1	1922	6313			
	Diethylene glycol dibenzoate		G	N <sup>i</sup>			34	1			0	0	
Bisphenols	Bisphenol A	BPA	L	T	50	1	38		461	1043	1	1	1
	Bisphenol S	BPS	L	T	55	1	38		691	5225	1		1
	Bisphenol A bis(2,3-dihydroxypropyl) ether	BADGE.2H <sub>2</sub> O	L	T	1000	1	16			11 237			
	Bisphenol A (3-chloro-2-hydroxypropyl) ether	BADGE-HCl-H <sub>2</sub> O	L	T	125	1	7			2205			
	Bisphenol AF	BPAF	L	S	1	1	3				1	1	
PBDEs	2,4,4'-Tribromodiphenyl ether	BDE-28	G	T	2.5	1	26		38	363			1
	2,2',4,4'-Tetrabromodiphenyl ether	BDE-47	G	T	1.0	1	38		720	6878			1
	2,2',4,4',5-Pentabromodiphenyl ether	BDE-99	G	T	5.0	1	36		721	5090			1
	2,2',4,4',5,5'-Hexabromodiphenyl ether	BDE-153	G	T	5.0	1	31		111	1033			1
	2,2',4,4',5,6'-Hexabromodiphenyl ether	BDE-154	G	T	2.5	1	31		88	872			1
	2,2',4,4',6-Pentabromodiphenyl ether	BDE-100	G	T	5.0	1	35		165	1394			1
	2,2',3,4,4',5,6-Heptabromodiphenyl ether	BDE-183	G	T	50	1	2			215			1

(Continues)

TABLE 1 (Continued)

Chemical class	Compound name	Abbreviation	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
OP-FRs	Tri-n-butyl phosphate	TNBP	G	T	100	1	19		100	2024	0	1	1
	Triphenyl phosphate	TPHP	G	T	5.0	1	37		2105	7120	1	1	1
	Tris(1-chloroisopropyl) phosphate	TCIPP	G	T	50	1	38		10 666	20 407	0		1
	Tris(4-butyl-phenyl) phosphate	TBPP	G	T	25	1	14			217			
	Tris(2-butoxyethyl) phosphate	TBOEP	G	T	250	1	37		7445	14 159	0		
	Tris(1,3-dichloro-2-propyl) phosphate	TDCPP	G	T	50	1	38		3955	15 338	1	1	1
	Tris(2-chloroethyl) phosphate	TCEP	G	T	25	1	37		2810	20 946	0	0	1
	Triethyl phosphate	TEP	L	S <sup>h</sup>	<100	1	8			154	0		
Other FRs	Tricresyl phosphate	TCP	L	S	25	1	8			8529	1		1
	Octyl diphenyl phosphate		L	S			3	1					
	Bis(2-ethylhexyl)-2,3,4,5-tetrabromophthalate	BEH-TEBP	G	T	1000	1	3			13 761	0		
	2,4-Dibromophenol		G	T	20	1	2			50			
	1,2-bis(2,4,6-tribromophenoxy) ethane	BTBPE	G	T	100	1	1						
	3,3',5,5'-Tetrabromobisphenol A	TBBPA	L	S <sup>h</sup>	<100	1	36		18	91	1	1	
	Melamine		L	S <sup>h</sup>	<100	1	17			3162			
	N-ethyl perfluorooctane sulfonamide ethanol	MeFOSE	L	T	8.0	1	6			90			1
PFAS	Perfluorobutane sulfonate	PFBS	L	T	2.0	1	9			23			1
	Perfluorohexane sulfonate	PFHxS	L	T	1.0	1	16			21			1
	Perfluorooctane sulfonate	PFOS	L	T	0.2	1	36		6	221	1	1	1
	Perfluorodecanoic acid	PFDA	L	T	4.0	1	21		11	97	1	0	1
	Perfluoroheptanoic acid	PFHpA	L	T	3.0	1	23		9	39	0	0	1
	Perfluorohexanoic acid	PFHxA	L	T	1.0	1	35		6	30	0	0	1
	Perfluorononanoic acid	PFNA	L	T	3.0	1	29		8	217	1	0	1
	Perfluorooctanoic acid	PFOA	L	T	3.0	1	34		10	138	0	0	1
	Perfluoropentanoic acid	PFPeA	L	T	3.0	1	12			10			1
	Perfluoroundecanoic acid	PFUnDA	L	S <sup>h</sup>	<100	1	5			125	0	0	1
	6:2 polyfluoroalkyl phosphate diester	6:2diPAP	L	N <sup>i</sup>		1	38						
	6:2diPAP/8:2diPAP		L	N <sup>i</sup>		1	38						
	Fluorotelomer sulfonic acid	FTSA 6:2	L	N <sup>i</sup>		1	38						
	1h-Perfluoroheptane		L	N			37	1					

(Continues)



TABLE 1 (Continued)

Chemical class	Compound name	Abbreviation	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
Phenols	2-Chlorophenol		G	T	50	1	1	1			1	0	
	2,6-Dichlorophenol		G	T	100	1	1	1					
	3,4,5-Trichlorophenol		G	T	500	1	1	1					
	Trichlorophenols (2,3,4-; 2,3,5-; 2,3,6-; 2,4,5-; 2,4,6-) <sup>j</sup>		G	T	100	1	1	1					
	Phenol		G	T	50	1	28		580	1669	1	0	
PAHs	Tetrachlorophenol		G	T	250	1	19	1		250			
	Cresol ( <i>o</i> -, <i>m</i> -, <i>p</i> -) <sup>i,k</sup>		G	T	100	1	34	1	250	250			
	2,4-Dinitrophenol		L	S <sup>j</sup>	50 000	1	7				0	1	
	Anthracene		G	T	5.0	1	6			24			
	Benzo(a)anthracene		G	T	5.0	1	10			89	1	1	1
	Benzo(g,h,i)perylene		G	T	50	1	17			289			
	Chrysene		G	T	5.0	1	18			130			1
	Phenanthrene		G	T	10	1	38		106	367	1	0	1
	Dibenzo(a,h)anthracene + Indeno(1,2,3-cd)pyrene <sup>l</sup>		G	T	50	1	5			260	0		
	Fluorene		G	T	10	1	1				0	0	1

Abbreviations: LOD, limit of detection; NHANES, National Health and Nutrition Examination Survey; OP-FRs, organophosphate flame retardants; PAHs, polycyclic aromatic hydrocarbons; PBDEs, polybrominated diphenyl ethers; PFAS, per- and polyfluoroalkyl substances.

<sup>a</sup>Type of analytical instruments used to detect compounds (G = gas chromatography and L = liquid chromatography).

<sup>b</sup>Type of analytical methods used to detect compounds (T = target analysis, S = suspect screening, and N = non-target analysis).

<sup>c</sup>Compounds confirmed by standards (1 = Yes and blank = No).

<sup>d</sup>Compounds newly detected in our dust (1 = Yes and blank = No).

<sup>e</sup>Compounds having endocrine-disrupting potential from in vitro high-throughput screening assays (1 = active, 0 = inactive, and blank = not tested or data not available).

<sup>f</sup>Compounds having neurotoxic potential from in vitro high-throughput screening assays (1 = active, 0 = inactive, and blank = not tested or data not available).

<sup>g</sup>Compounds biomonitoring in NHANES (1 = Yes and blank = No).

<sup>h</sup>Detected by suspect screening or non-target methods first and quantified with standards later, thus concentrations are semi-quantitative.

<sup>i</sup>Detected by suspect screening or non-target methods first and quantified with standards later, but concentrations were not quantified.

<sup>j</sup>Detected with multiple isomers.

<sup>k</sup>Detected in most samples (above LOD) but were below the limit of quantification (LOQ).

<sup>l</sup>Structures of these compounds were too close to discriminate one from the other.



(DEP) and dimethyl phthalate (DMP) were detected in 79% and 71% of the samples, respectively, at relatively low concentrations (medians were below 1000 ng/g of dust). Di-n-octyl phthalate (DOP), a target compound of the current study, was not detected in our dust, whereas it was detected in 100% of other California house dust samples collected in 2006.<sup>2</sup>

We detected 6 non-phthalate plasticizers and four were newly detected in our dust (dioctyl terephthalate (DOTP), 1,3-diphenylguanidine, toluene-2-sulfonamide, and diethylene glycol dibenzoate). Note that 1,3-diphenylguanidine is primarily used in various solid items including rubber footwear and automobile tires. Four out of six compounds were widely detected in our samples ( $\geq 34$  out of 38 samples). The latter two in Table 1 were detected via suspect screening and non-target methods, respectively. The median concentration of DOTP, a direct replacement for DEHP, was as high as DEHP. Among the four newly detected compounds, 1,3-diphenylguanidine has both endocrine and neurotoxic potential, but has not been biomonitoring in NHANES.

### 3.2.2 | Bisphenols and bisphenol analogs (hereafter referred as “bisphenols”)

BPA and BPS were detected in all of our samples, and two bisphenol analogs that can serve as replacements for BPA were measured in our dust samples via a target method, including bisphenol A bis(2,3-dihydroxypropyl) ether [BADGE-2H<sub>2</sub>O] and bisphenol A (3-chloro-2-hydroxypropyl) (2,3-dihydroxypropyl) ether [BADGE-HCl-H<sub>2</sub>O]. Bisphenol AF (BPAF) was also detected in our samples via a suspect screening method. We confirmed that BPAF has both endocrine and neurotoxic potential, but has not been biomonitoring in NHANES. BADGE-2H<sub>2</sub>O and BADGE-HCl-H<sub>2</sub>O were not tested for endocrine-disrupting and neurotoxic potential using in vitro HTS assays and have not been biomonitoring in NHANES.

### 3.2.3 | PBDEs, OP-FRs, and other FRs

Seven PBDEs were detected and quantified in our samples via a target method, and they have been biomonitoring in NHANES. Although in vitro toxicity screening data were not available for all PBDEs, other adverse health effects of PBDEs have been summarized elsewhere.<sup>45</sup> BDE-209 has been widely detected in other California house dust.<sup>46,47</sup> However, we could not have detected BDE-209 in our samples because our GC method was not designed to measure compounds with such low volatility.

Ten OP-FRs, including 7 target compounds, were detected in our dust samples. Five OP-FRs were ubiquitous ( $>97\%$ ) in our samples. Overall, median concentrations of OP-FRs were higher than those of PBDEs by one order of magnitude. Even though tris(2-butoxyethyl) phosphate (TBOEP) was ubiquitous ( $>97\%$ ) in our samples and measured at high concentrations (median was 7445 ng/g of dust), it has not been biomonitoring in NHANES. Octyl diphenyl phosphate was newly detected in our dust via

suspect screening. Four OP-FRs have either endocrine-disrupting or neurotoxic potential, but in vitro toxicity screening data were not available for tris(4-butyl-phenyl) phosphate (TBPP) and octyl diphenyl phosphate.

In addition to PBDEs and OP-FRs, we detected five compounds that are used as flame retardants. Three compounds were detected in a few samples ( $<4$ ) via a target method. Melamine and 3,3',5,5'-Tetrabromobisphenol A (TBBPA) were detected in 17 and 36 samples, respectively, via suspect screening. TBBPA has both endocrine-disrupting potential and neurotoxic potential, but has not been biomonitoring in NHANES. Because melamine was detected in almost half of our samples and has been detected in urine specimens of children who consumed milk products,<sup>48</sup> toxicity testing and biomonitoring are recommended for this compound.

### 3.2.4 | PFAS

A total of 15 PFAS were detected in our dust, including 10 target compounds. We newly detected 3-(perfluorooctyl)propyl iodide in our dust (37 out of 38 samples) via a non-target method. Compared with other chemical classes, PFAS median concentrations were relatively low ( $<12$  ng/g of dust). Eleven PFAS have been biomonitoring in NHANES and seven of them were tested for endocrine-disrupting and neurotoxic potential using in vitro HTS assays.

### 3.2.5 | Phenols

Among 15 targeted phenols, only 7 phenols were detected in our samples whereas trichlorophenols and cresols were detected with multiple isomers. Four phenols were newly detected in our dust via a target method, but in only one sample. We additionally detected 2,4-dinitrophenol via suspect screening. Overall, due to the low detection frequency, median concentrations were computed only for phenol and cresols (580 and 250 ng/g of dust, respectively). None of the 8 detected phenols was biomonitoring in NHANES, and only three phenols were tested for toxicity using in vitro HTS assays. Cresols were detected with multiple isomers (o-, m-, and p-) in 34 out of 38 samples and thus are recommended to be included in future biomonitoring and in vitro toxicity screening testing.

### 3.2.6 | PAHs

Among 12 targeted PAHs, 8 PAHs were detected in our samples. Because structures of dibenzo(a,h)anthracene and indeno(1,2,3-cd)pyrene were too close to discriminate one from the other, results were reported together in Table 1. Except for phenanthrene, 7 other PAHs were detected in fewer than 50% of our samples. Four PAHs were or have been biomonitoring in NHANES. In vitro toxicity screening testing data for endocrine-disrupting potential or neurotoxic potential were not available for three PAHs, but other toxic endpoints are available from in vitro HTS assays.<sup>49</sup>

TABLE 2 Summary of biocides that were detected in our dust samples (n = 38)

Chemical class	Compound name	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
Insecticides	Fipronil-sulfone	L	T	1.0	1	37		77	3004			
	Fipronil	L	T	1.0	1	37		62	5718	1	1	
	Permethrin	G	T	50	1	34		1922	8151	1	0	1
	Imidacloprid	L	T	10	1	32		123	22 455	0	0	1
	Bifenthrin	G	T	5.0	1	31		140	3336	1	0	
	Fipronil-desulfinyl	L	T	2.0	1	28		11	248			
	Fipronil-sulfide	L	T	1.0	1	17			136			
	Novaluron	L	T	1.0	1	9	1		1826	1	0	
	Chlorpyrifos	G	T	5.0	1	8			365	1	0	1
	Methoxyfenozide	L	T	2.0	1	7	1		204	1	0	
	Propoxur	L	T	6.0	1	6			60	0	0	1
	Chlorantraniprole	L	T	3.0	1	4	1		69			
	Pyriproxyfen	L	T	4.0	1	4			791	0	1	
	Deltamethrin	G	T	500	1	4			6658	0		1
	Etofenprox	G	T	10	1	4			38 059	1	0	
	Cyhalothrin	G	T	50	1	2			7449	0		1
	Cypermethrin	G	T	500	1	2			24 222	1		1
	1,3-benzothiazole	L	T	600	1	2	1		2714	0		
	Esfenvalerate	G	T	250	1	1				0	1	
	Tetrachlorvinphos	L	S <sup>h</sup>	<100	1	6			43 222	1		1
	Cyfluthrin	G	N <sup>i</sup>		1	1						1
	Piperonyl butoxide	G	N			26				1	0	

(Continues)

TABLE 2 (Continued)

Chemical class	Compound name	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
Fungicides	Boscalid	L	T	3.0	1	10			262	0	1	
	Azoxystrobin	L	T	5.0	1	11	1		77	1	1	
	Difenoconazole	L	T	9.0	1	2	1		18	1	1	
	Didecylidimethylammonium chloride	L	T	100	1	28	1	2859	10 579	1	1	
	Pentachlorophenol	G	T	1000	1	6			11 511	1	0	1
	Carbendazim	L	S <sup>h</sup>	<100	1	11			2568	1	0	
	Imazalil	L	S <sup>h</sup>	<100	1	29		18	210	1	1	
	Octhilinone	L	S <sup>h</sup>	<100	1	22		5	44	1	1	
	Propiconazole	L	S <sup>h</sup>	<100	1	31		34	194	1	1	
	Thiabendazole	L	S <sup>h</sup>	<100	1	37		54	713	0	0	
	Fludioxonil	L	S <sup>h</sup>	<100	1	37	1	12	61	1	1	
	Physcion	L	S <sup>h</sup>	<100	1	38		177	866			
	4-hydroxychlorothalonil	L	N <sup>i</sup>		1	38	1					
	Pyraclostrobin	L	S <sup>j</sup>		1	7				0	1	
	Dichlorophen	L	S	250	1	1	1			1		
Herbicides	3-Iodo-2-propynyl-N-butylcarbamate	L	S	50	1	16	1		385	1	1	
	2,4-Dichlorophenoxyacetic acid	L	T	100	1	20		500	5057	0	0	1
	Propanil	L	T	2.0	1	13			512	1	1	
	Diuron	L	T	15	1	11			1116	1	0	
	Pendimethalin	L	T	150	1	5			591	1	1	
	Simazine	L	T	3.0	1	1				0	0	

Note: For a through i, refer to footnote of Table 1.

Abbreviations: LOD, limit of detection; NHANES, National Health and Nutrition Examination Survey.

### 3.3 | Biocides

#### 3.3.1 | Insecticides

We detected 22 insecticides mostly via a target method (Table 2). Four insecticides were newly detected in our samples, but in fewer than 25% of the samples. Permethrin was measured at the highest median concentration (1922 ng/g of dust), but the 95th percentile concentration of other four insecticides (imidacloprid, etofenprox, cypermethrin, and tetrachlorvinphos) was higher than that of permethrin. In addition to common insecticides used to control pests indoors, other potential indoor sources of fipronil, imidacloprid, pyriproxyfen and permethrin may be associated with their use as topical flea control agents for dogs and cats. About 53% of the participating homes had at least one indoor cat or dog. Fipronil products have been shown to persist on pets for over 28 days,<sup>50</sup> and this is likely true for other active ingredients based on the relatively low frequency of application required for these products. These active ingredients are likely to accumulate in dust as pets shed treated fur and skin cells. The less frequent detection of some of the insecticides in this study, coupled with their relatively high coefficients of variation (see Table S2), likely reflects the fact that pet ownership and indoor insecticide applications are not as ubiquitous as other indoor product uses. Compared with toxicity testing data ( $n = 16$ ), biomonitoring data are limited ( $n = 9$ ). For fipronil-sulfone, fipronil, fipronil-desulfinyl, and fipronil-sulfide that were detected in around or greater than 50% of the samples, both toxicity testing and biomonitoring are recommended.

#### 3.3.2 | Fungicides

A total of 15 fungicides were detected, 11 of them via suspect screening or non-target methods. Seven fungicides were newly detected in our dust. Eight fungicides mostly detected via suspect screening or non-target methods had a detection frequency above 50%, indicating widespread use of fungicides in the indoor environment. Except for didecyldimethylammonium chloride (DDAC; median concentration = 2859 ng/g of dust), measured median concentrations of fungicides were low compared with insecticides. In addition to a fungicidal use, DDAC is used as an antibacterial agent and has wide indoor applications where it is used on walls, floors, tables, toilets, and fixtures.<sup>51</sup> Of the 15 detected fungicides, 13 fungicides were previously tested for either endocrine-disrupting potential or neurotoxic potential. Except for pentachlorophenol, none of the 14 detected fungicides has been biomonitoring in NHANES. Among the compounds that were newly detected and have both endocrine-disrupting and neurotoxic potential, two fungicides (DDAC and fludioxonil) were commonly detected (>70%) in our samples; two other fungicides (azoxystrobin and difenoconazole) were detected infrequently (29% and 5%, respectively). Thus, fungicides with a high detection frequency are recommended to be included in future biomonitoring studies.

#### 3.3.3 | Herbicides

Five herbicides were detected in our dust via a target method. The detection frequency was below 50% for 4 out of 5 herbicides. Given the low detection frequency, they might be attributable to applications in agricultural fields or gardens. Except for 2,4-dichlorophenoxyacetic acid, none of them were biomonitoring in NHANES. Propanil and pendimethalin showed both endocrine-disrupting potential and neurotoxic potential, and diuron showed endocrine-disrupting potential from the *in vitro* testing. Thus, they might need to be included in future biomonitoring studies.

### 3.4 | Compounds in PCPs

We detected parabens, fragrance ingredients, UV filters, cosmetic ingredients, and those with other personal care uses in our dust (Table 3). Overall, they were widely detected, and median concentrations were on the order of 1000–10 000 ng/g of dust for 11 compounds. For users of products containing these compounds, direct dermal uptake is likely to be a primary exposure route. However, for non-users, such as young children who spend most of their time on the floors and have high dust ingestion rates, dust may be an important exposure medium for these compounds.<sup>52</sup> Eight compounds in PCPs were newly detected in our dust and four compounds are cosmetic ingredients. Among newly detected compounds, toxicity testing and biomonitoring are recommended for dexpanthenol because it was detected in all samples with a median of 1311 ng/g of dust. There are only 8 compounds that have been biomonitoring in NHANES and that were tested for endocrine-disrupting and/or neurotoxic potential. However, we observed that there are many PCP compounds that may require biomonitoring and toxicity testing based on the detection frequency and high median concentrations.

### 3.5 | Chemical classes whose dust concentrations are of less concern for environmental exposure calculations

In our samples, we detected 10 food additives, 5 sweeteners, 11 food sources, 29 pharmaceuticals, 3 skin oils, 14 human metabolites, 31 surfactants, and 17 compounds whose use is not known (Table S2). Most of them were detected via suspect screening or non-target methods. These compound classes were not of interest in identifying or measuring dust concentrations in previous indoor environmental monitoring studies. Thus, their presence was rarely reported in the literature and 87 out of 119 newly detected compounds fell in this category. Among the compounds classified in this category, only 18 compounds were tested for endocrine-disrupting and/or neurotoxic potential. We found that sorbic acid (food additive), ketoconazole (pharmaceutical), nicotine (pharmaceutical), linoleic acid (food sources), linolenic acid (food sources), and genistein (food sources) have endocrine-disrupting and/or neurotoxic potential. Exposure to genistein occurs primarily through foods made with soybeans and

**TABLE 3** Summary of compound classes that were detected in our dust samples (n = 38) and mainly used in dermal applications

Chemical class	Compound name	Instrument <sup>a</sup>	Method <sup>b</sup>	LOD (ng/g of dust)	Standard <sup>c</sup>	# detections	New detection <sup>d</sup>	Median conc. (ng/g of dust)	95th percentile (ng/g of dust)	Endocrine <sup>e</sup>	Neurotoxic <sup>f</sup>	NHANES <sup>g</sup>
Parabens	Ethyl paraben	L	T	2.0	1	35		48	241	1	1	1
	Methyl paraben	L	T	6.0	1	37		524	1559	0	0	1
	Butyl Paraben + Isobutyl Paraben <sup>i</sup>	L	T	2.0	1	26		23	235	1	1	1
	Isopropyl paraben + Propyl Paraben <sup>i</sup>	L	T	2.0	1	38		226	1457	1	1	1
	2-Benzylideneoctanal	G	T	100	1	37	1	1194	3950	0	0	
Fragrance	Galaxolide	G	T	5.0	1	38		1294	5521			
	Tonalide	G	T	10	1	27		219	818	0		
	Homosalate	G	T	25	1	38		5233	17 398	0		
UV filters	Octocrylene	G	T	25	1	38		2245	13 656			
	2,4-dihydroxybenzophenone	L	T	1.0	1	35		22	148	1		
	Benzophenone-4	L	S <sup>h</sup>	<100	1	30		37	230			
	Octyl methoxycinnamate	L	S <sup>h</sup>	<100	1	37		1450	8037			
	2-Ethylhexyl salicylate	G	N <sup>h</sup>	n.d.	1	38		7057	26 531			
	Benzophenone	G	N <sup>h</sup>	n.d.	1	35		441	1326	0	0	
	Benzophenone-3	G	N <sup>h</sup>	n.d.	1	38		19 170	64 114	0		1
	Triclocarban	L	T	1.0	1	37		33	1144	1	1	1
	Triclosan	L	T	5.0	1	38		275	1391	1	1	1
	N,N-Diethyl-meta-tolamide (DEET)	L	T	15	1	30		154	5536	1	0	1
Other personal care	Dexpantenol	L	S <sup>h</sup>	<100	1	38	1	1311	13 243			
	Benzyl Benzoate	G	N <sup>h</sup>	n.d.	1	38	1	1280	10 628	0		
	Salnacedin	L	N			38	1					
	Coumarin	G	N			33				0	0	
	Ethyl N-acetyl-L-N-butyl-β-alaninate	L	S			4						
	Tricaprylin	G	N <sup>h</sup>	n.d.	1	33		27 882	75 596			
	Isopropyl myristate	G	N <sup>h</sup>	n.d.	1	38	1	1836	8485	0		
Cosmetics	Lilial	G	N <sup>h</sup>	n.d.	1	38	1	319	3557	0		
	Cholesteryl benzoate	G	N			38	1					
	Isopropyl palmitate	G	N			38						
	12-Hydroxystearic acid	L	N			38	1					
	Benzyl salicylate	G	N			38				0		

Note: For a thorough h, refer to footnote of Table 1.  
Abbreviations: n.d., not determined.  
<sup>i</sup>Structures of these compounds were too close to discriminate one from the other.

soy protein.<sup>53</sup> Biomonitoring is recommended for linoleic acid because it was detected in all samples at a high median concentration (34 308 ng/g of dust) and has endocrine-disrupting potential.

## 4 | DISCUSSION

Results from this study provided more comprehensive chemical profiles of house dust. We detected a total of 276 compounds in our dust samples and quantified concentrations of 144 compounds using standards. In addition to the compounds that were previously measured in indoor dust, we tried to identify overlooked compounds that were not previously measured in dust but were shown to have potential for adverse health effects from the HTS toxicity testing. We were also able to expand the list of compounds present in indoor dust by applying both LC-MS and GC-MS with three analytical approaches. For example, 75% of the newly measured chemicals were observed via LC non-target or LC suspect approaches (see Figure S1). The newly measured compounds in our study mainly comprised surfactants, pharmaceuticals, and human metabolites. Because of the polarity of these compounds, we were able to detect a large number of compounds via LC-MS. Another reason we could extend the list of compounds present in indoor dust is that our samples were recently collected (2015-2016). Compared with Rager et al.<sup>22</sup> who investigated non-targeted compounds using LC-MS in U.S. household dust samples collected from 2005 to 2006, we were able to newly detect four replacement plasticizers (eg, acetyl tributyl citrate and DOTP) in our samples, reflecting currently used products.

Our study showed that indoor dust contains chemicals from various consumer product uses and also supported the idea that dust can serve as a marker of use. For example, most of the food additives and sweeteners detected in our dust are used in processed foods or drinks. Thus, the presence of food additives or sweeteners in dust indicates that they exist outside their intended use, which is to be consumed via direct food intake. Cholesterol (found in skin and emitted during cooking) and skin oils were ubiquitously measured in Danish homes and daycare centers.<sup>44</sup> In addition to these compounds, we observed cosmetic ingredients and vitamin E with median concentrations greater than 10 000 ng/g. In a separate study in which we analyzed skin wipe samples,<sup>54</sup> 11 compounds (triethyl citrate, butylated hydroxytoluene, cholesta-3,5-diene, vitamin E, cholesterol, tridecanoic acid, arachidonic acid, palmitol, palmitic acid, pentadecanoic acid, and linolenic acid) were detected, and they were also detected in our dust samples. This indicates that human activities, including cooking, cosmetic use, skin sloughing, dropping food residue, or debris unintentionally on floors, could be sources of these compounds. Moreover, because we analyzed recently collected dust samples, we observed that relatively new chemicals (eg, OP-FRs and BPS) were measured at higher concentrations than those for controversial or banned chemicals in consumer products (eg, PBDEs and BPA). This reflects the dynamic nature of consumer product formulations, especially given heightened consumer awareness and concerns about the safety of product ingredients.

Compiling existing exposure and toxicity potential data of our detected compounds allowed us to inform key data gaps for assessing potential health effects for previously overlooked chemicals. For example, we found that in vitro HTS toxicity data were not available for some of the detected plasticizers, bisphenols, and biocides, which may adversely affect human health. Of most interest are one plasticizer (toluene-2-sulfonamide), two bisphenols (BADGE.2H<sub>2</sub>O and BADGE-HCl-H<sub>2</sub>O), and eight biocides and biocide transformation products (fipronil-desulfinyl, fipronil-sulfide, fipronil-sulfone, chlorantraniprole, cypermethrin, cyfluthrin, 4-hydroxychlorothalonil, and physcion). Because most of these compounds were ubiquitous in our samples and may have toxicity potential, they are recommended to be included in future in vitro toxicity screening.

In conclusion, following the identification of a broad spectrum of chemicals from a previous study,<sup>18</sup> this study integrated their measured dust concentrations with existing exposure and toxicity information to inform key data gaps for assessing potential health effects for consumer product chemicals. We found that 13 newly detected compounds may potentially disrupt endocrine systems and/or be neurotoxic based on in vitro bioactivity assays. These results expand our knowledge of chemicals present in indoor residential environments where vulnerable populations, especially young children, spend most of their time on the floors.<sup>55</sup> Consequently, we expect that our findings may trigger further environmental health research regarding previously overlooked compounds. Many of the pharmaceuticals and PCPs newly detected in our dust have been extensively studied in various aquatic environments, including drinking water, wastewater, surface water, and groundwater<sup>56</sup> because they may pose a threat to the ecosystem and/or human health. Given that people spend most of their time indoors,<sup>55</sup> more studies are needed to examine the presence of these compounds in residential dust and to investigate potential health effects associated with indoor non-dietary exposure routes. Additional studies are also recommended to confirm the presence of compounds that were less frequently detected in the current study and not yet confirmed by standards.

## ACKNOWLEDGEMENTS

Research reported in this publication was supported by the US Environmental Protection Agency (EPA-G2013-STAR-K1) and the UC Davis Superfund Research Center, National Institutes of Health, NIEHS award (P42-ES004699). All authors declare they have no actual or potential competing financial interests.

## CONFLICT OF INTEREST

The authors declare that they have no competing interest.

## ORCID

Hyeong-Moo Shin  <https://orcid.org/0000-0002-7001-2555>

## REFERENCES

- Egeghy PP, Vallero DA, Hubal E. Exposure-based prioritization of chemicals for risk assessment. *Environ Sci Policy*. 2011;14(8):950-964.
- Dodson RE, Camann DE, Morello-Frosch R, Brody JG, Rudel RA. Semivolatile organic compounds in homes: strategies for efficient and systematic exposure measurement based on empirical and theoretical factors. *Environ Sci Technol*. 2015;49(1):113-122.
- Mitro SD, Dodson RE, Singla V, et al. Consumer product chemicals in indoor dust: a quantitative meta-analysis of US studies. *Environ Sci Technol*. 2016;50(19):10661-10672.
- Zota AR, Calafat AM, Woodruff TJ. Temporal trends in phthalate exposures: findings from the national health and nutrition examination survey, 2001-2010. *Environ Health Persp*. 2014;122(3):235-241.
- Calafat AM, Valentin-Blasini L, Ye X. Trends in exposure to chemicals in personal care and consumer products. *Curr Environ Health Rep*. 2015;2(4):348-355.
- Cordner A, Mulcahy M, Brown P. Chemical regulation on fire: rapid policy advances on flame retardants. *Environ Sci Technol*. 2013;47(13):7067-7076.
- Shin HM, Ernstoff A, Arnot JA, et al. Risk-based high-throughput chemical screening and prioritization using exposure models and in vitro bioactivity assays. *Environ Sci Technol*. 2015;49(11):6760-6771.
- Xu Y, Hubal E, Little JC. Predicting residential exposure to phthalate plasticizer emitted from vinyl flooring: sensitivity, uncertainty, and implications for biomonitoring. *Environ Health Persp*. 2010;118(2):253-258.
- Beko G, Weschler CJ, Langer S, Callesen M, Toftum J, Clausen G. Children's phthalate intakes and resultant cumulative exposures estimated from urine compared with estimates from dust ingestion, inhalation and dermal absorption in their homes and daycare centers. *PLoS ONE*. 2013;8(4):e62442.
- Little JC, Weschler CJ, Nazaroff WW, Liu Z, Hubal E. Rapid methods to estimate potential exposure to semivolatile organic compounds in the indoor environment. *Environ Sci Technol*. 2012;46(20):11171-11178.
- Shin HM, McKone TE, Tulse NS, Clifton MS, Bennett DH. Indoor residence times of semivolatile organic compounds: model estimation and field evaluation. *Environ Sci Technol*. 2013;47(2):859-867.
- Weschler CJ, Nazaroff WW. SVOC partitioning between the gas phase and settled dust indoors. *Atmos Environ*. 2010;44(30):3609-3620.
- Ao JJ, Yuan T, Ma YN, Gao L, Ni N, Li D. Identification, characteristics and human exposure assessments of triclosan, bisphenol-A, and four commonly used organic UV filters in indoor dust collected from Shanghai, China. *Chemosphere*. 2017;184:575-583.
- Hwang HM, Park EK, Young TM, Hammock BD. Occurrence of endocrine-disrupting chemicals in indoor dust. *Sci Total Environ*. 2008;404(1):26-35.
- Lankova D, Svarcova A, Kalachova K, Lacina O, Pulkrabova J, Hajslova J. Multi-analyte method for the analysis of various organohalogen compounds in house dust. *Anal Chim Acta*. 2015;854:61-69.
- Dodson RE, Perovich LJ, Covaci A, et al. After the PBDE phase-out: a broad suite of flame retardants in repeat house dust samples from California. *Environ Sci Technol*. 2012;46(24):13056-13066.
- Rudel RA, Camann DE, Spengler JD, Korn LR, Brody JG. Phthalates, alkylphenols, pesticides, polybrominated diphenyl ethers, and other endocrine-disrupting compounds in indoor air and dust. *Environ Sci Technol*. 2003;37(20):4543-4553.
- Moschet C, Anumol T, Lew BM, Bennett DH, Young TM. Household dust as a repository of chemical accumulation: new insights from a comprehensive high-resolution mass spectrometry study. *Environ Sci Technol*. 2018;52:2878-2887.
- Krauss M, Singer H, Hollender J. LC-high resolution MS in environmental analysis: from target screening to the identification of unknowns. *Anal Bioanal Chem*. 2010;397(3):943-951.
- Hilton DC, Jones RS, Sjodin A. A method for rapid, non-targeted screening for environmental contaminants in household dust. *J Chromatogr A*. 2010;1217(44):6851-6856.
- Ouyang XY, Weiss JM, de Boer J, Lamoree MH, Leonards P. Non-target analysis of household dust and laundry dryer lint using comprehensive two-dimensional liquid chromatography coupled with time-of-flight mass spectrometry. *Chemosphere*. 2017;166:431-437.
- Rager JE, Strynar MJ, Liang S, et al. Linking high resolution mass spectrometry data with exposure and toxicity forecasts to advance high-throughput environmental monitoring. *Environ Int*. 2016;88:269-280.
- Rostkowski P, Haglund P, Aalizadeh R, et al. The strength in numbers: comprehensive characterization of house dust using complementary mass spectrometric techniques. *Anal Bioanal Chem*. 2019;411(10):1957-1977.
- Judson RS, Houck KA, Kavlock RJ, et al. In vitro screening of environmental chemicals for targeted testing prioritization: the ToxCast project. *Environ Health Perspect*. 2010;118(4):485-492.
- Isaacs KK, Glen WG, Egeghy P, et al. SHEDS-HT: an integrated probabilistic exposure model for prioritizing exposures to chemicals with near-field and dietary sources. *Environ Sci Technol*. 2014;48(21):12750-12759.
- Shin HM, McKone TE, Bennett DH. Intake fraction for the indoor environment: a tool for prioritizing indoor chemical sources. *Environ Sci Technol*. 2012;46(18):10063-10072.
- Zhang X, Arnot JA, Wania F. Model for screening-level assessment of near-field human exposure to neutral organic chemicals released indoors. *Environ Sci Technol*. 2014;48(20):12312-12319.
- Wenger Y, Li D, Jolliet O. Indoor intake fraction considering surface sorption of air organic compounds for life cycle assessment. *Int J Life Cycle Assess*. 2012;17(7):919-931.
- Centers for Disease Control and Prevention. *Fourth National Report on Human Exposure to Environmental Chemicals*. Atlanta, GA: Centers for Disease Control and Prevention; 2019:1-235.
- Blanchard O, Glorennec P, Mercier F, et al. Semivolatile organic compounds in indoor air and settled dust in 30 French dwellings. *Environ Sci Technol*. 2014;48(7):3959-3969.
- Pham N, Iyer S, Hackett E, et al. Using ToxCast to explore chemical activities and hazard traits: a case study with ortho-phthalates. *Toxicol Sci*. 2016;151(2):286-301.
- Roberts JW, Budd WT, Ruby MG, et al. Development and field testing of a high volume sampler for pesticides and toxics in dust. *J Expo Anal Environ Epidemiol*. 1991;1(2):143-155.
- Goldsmith MR, Grulke CM, Brooks RD, et al. Development of a consumer product ingredient database for chemical exposure screening and prioritization. *Food Chem Toxicol*. 2014;65:269-279.
- He LY, Hu M, Huang XF, Yu BD, Zhang YH, Liu DQ. Measurement of emissions of fine particulate organic matter from Chinese cooking. *Atmos Environ*. 2004;38(38):6557-6564.
- Dodson RE, Nishioka M, Standley LJ, Perovich LJ, Brody JG, Rudel RA. Endocrine disruptors and asthma-associated chemicals in consumer products. *Environ Health Persp*. 2012;120(7):935-943.
- Smith MN, Grice J, Cullen A, Faustman EM. A toxicological framework for the prioritization of children's safe product act data. *Int J Environ Res Public Health*. 2016;13(4):431.
- Kleinstreuer NC, Ceger P, Watt ED, et al. Development and validation of a computational model for androgen receptor activity. *Chem Res Toxicol*. 2017;30(4):946-964.
- Rotroff DM, Martin MT, Dix DJ, et al. Predictive endocrine testing in the 21st century using in vitro assays of estrogen receptor signaling responses. *Environ Sci Technol*. 2014;48(15):8706-8716.



39. Friedman KP, Watt ED, Hornung MW, et al. Tiered high-throughput screening approach to identify thyroperoxidase inhibitors within the toxcast Phase I and II chemical libraries. *Toxicol Sci.* 2016;151(1):160-180.
40. Rotroff DM, Dix DJ, Houck KA, et al. Using in vitro high throughput screening assays to identify potential endocrine-disrupting chemicals. *Environ Health Persp.* 2013;121(1):7-14.
41. Karmaus AL, Toole CM, Filer DL, Lewis KC, Martin MT. High-throughput screening of chemical effects on steroidogenesis using H295R human adrenocortical carcinoma cells. *Toxicol Sci.* 2016;150(2):323-332.
42. Strickland JD, Martin MT, Richard AM, Houck KA, Shafer TJ. Screening the ToxCast phase II libraries for alterations in network function using cortical neurons grown on multi-well microelectrode array (mwMEA) plates. *Arch Toxicol.* 2018;92(1):487-500.
43. Hornung RW, Reed LD. Estimation of average concentration in the presence of nondetectable values. *Appl Occup Environ Hyg.* 1990;5(1):46-51.
44. Weschler CJ, Langer S, Fischer A, Beko G, Toftum J, Clausen G. Squalene and cholesterol in dust from Danish homes and daycare centers. *Environ Sci Technol.* 2011;45(9):3872-3879.
45. Agency for Toxic Substances and Disease Registry. *Toxicological Profile for Polybrominated Diphenyl Ethers (PBDEs)*. Atlanta, GA: Agency for Toxic Substances and Disease Registry; 2017.
46. Guo WH, Park JS, Wang YZ, et al. High polybrominated diphenyl ether levels in California house cats: House dust a primary source? *Environ Toxicol Chem.* 2012;31(2):301-306.
47. Shen B, Whitehead TP, McNeel S, et al. High levels of polybrominated diphenyl ethers in vacuum cleaner dust from California fire stations. *Environ Sci Technol.* 2015;49(8):4988-4994.
48. Gabriels G, Lambert M, Smith P, Wiesner L, Hiss D. Melamine contamination in nutritional supplements – is it an alarm bell for the general consumer, athletes, and 'Weekend Warriors'? *Nutr J.* 2015;14. <https://nutritionj.biomedcentral.com/track/pdf/10.1186/s12937-015-0055-7>
49. Williams AJ, Grulke CM, Edwards J, et al. The CompTox Chemistry Dashboard: a community data resource for environmental chemistry. *J Cheminformatics.* 2017;9. <https://jcheminf.biomedcentral.com/track/pdf/10.1186/s13321-017-0247-6>
50. Teerlink J, Hernandez J, Budd R. Fipronil washoff to municipal wastewater from dogs treated with spot-on products. *Sci Total Environ.* 2017;599:960-966.
51. National Center for Biotechnology Information. *PubChem Database*. <https://pubchem.ncbi.nlm.nih.gov/compound/Didecyltrimethylammonium-chloride#section=Overview>. Accessed July 29, 2019.
52. U.S. Environmental Protection Agency. *Exposure Factors Handbook*. Washington, DC: U.S. Environmental Protection Agency; 2011.
53. Barlow J, Johnson JAP, Scofield L. *Early Life Exposure to the Phytoestrogen Genistein and Breast Cancer Risk in Later Years: Fact Sheet – Phytoestrogen Genistein*. Madison, Wisconsin: Breast Cancer & The Environment Research Centers; 2007.
54. Alfonso-Garrido J, Bennett DH, Parthasarathy S, Moschet C, Young TM, McKone TE. Exposure assessment for air-to-skin uptake of semivolatile organic compounds (SVOCs) indoors. *Environ Sci Technol.* 2019;53(3):1608-1616.
55. Klepeis NE, Nelson WC, Ott WR, et al. The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants. *J Expo Anal Environ Epidemiol.* 2001;11(3):231-252.
56. Ebele AJ, Abou-Elwafa Abdallah M, Harrad S. Pharmaceuticals and personal care products (PPCPs) in the freshwater aquatic environment. *Emerg Contam.* 2017;3(1):1-16.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Shin H-M, Moschet C, Young TM, Bennett DH. Measured concentrations of consumer product chemicals in California house dust: Implications for sources, exposure, and toxicity potential. *Indoor Air.* 2020;30:60-75. <https://doi.org/10.1111/ina.12607>